arXiv:1209.4396v2 [math.NT] 28 Sep 2012

CHEBYSHEV ACTION ON FINITE FIELDS

THOMAS ALDEN GASSERT

Abstract. Given a polynomial $\phi(x)$ and a finite field $\mathbb{F}_{p^n}$ one can construct a directed graph where the vertices are the values in the finite field, and emanating from each vertex is an edge joining the vertex to its image under ϕ. When ϕ is a Chebyshev polynomial of prime degree, the graphs display an unusual degree of symmetry. In this paper we provide a complete description of these graphs, and also provide some examples of how these graphs can be used to determine the decomposition of primes in certain field extensions.

1. Introduction

Let K be a number field and let $\phi \in K[x]$ a polynomial of degree at least 2. Define $\phi^{\circ n}(x) = \phi(\phi^{\circ(n-1)}(x))$ to be the n-fold iterate of $\phi(x)$ under composition with $\phi^{\circ 0}(x) = x$.

Let S be a set that accepts the action from ϕ. We say that $s \in S$ is periodic with period $c \geq 1$ if c is the minimal positive integer for which $\phi^{\circ c}(s) = s$. We say that $s \in S$ is preperiodic with preperiod $t \geq 0$ if t is the minimal non-negative integer such that $\phi^{\circ t}(s)$ is periodic. It is a simple conclusion that if S is a finite set, then every $s \in S$ is preperiodic.

Given ϕ and S, we can construct a directed graph with vertices being the elements of s and edges $(s \to \phi(s))$. In general, the structure of these graphs can be difficult to describe. However, when ϕ is taken to be a Chebyshev polynomial and S a finite field, the resulting graphs display an unusual degree of symmetry which would otherwise be absent for more general choices of ϕ. Furthermore, the structure of these graphs are both predictable and visually pleasing. The case of the Chebyshev polynomial $x^2 - 2$ acting on $\mathbb{F}_p$ was previously studied by T. Vasiga and J. Shallit in their paper [3]. Section 2 of our paper generalizes their results to a complete description of the graphs of Chebyshev polynomials of prime degree acting on arbitrary finite fields $\mathbb{F}_{p^n}$.

One motivation for studying these graphs is a paper by W. Aitken, F. Hajir, and C. Maire [1] which describes, under specific conditions, a method of using graphs to determine how unramified primes decompose in iterated field extensions. To understand their result, we provide some additional definitions.

Notation 1. Let $\phi \in K[x]$ as before. We call $\mathcal{R}_\phi = \{r \in \overline{K} \colon \phi'(r) = 0\}$ the critical points of ϕ, and $\mathcal{B}_\phi = \{\phi(r) \colon r \in \mathcal{R}_\phi\}$ the critical values. We say that ϕ is postcritically finite if every critical point is preperiodic for ϕ. Lastly, we define a family of polynomials

$$\Phi_n(x,t) = \phi^{\circ n}(x) - t \in K(x)[t].$$

Proposition 1.1. [Aitken, Hajir, Maire] Suppose that $\phi(x) \in \mathcal{O}_K[x]$ is postcritically finite and that $t \in \mathcal{O}_K$ is such that $\Phi_n(x,t)$ is irreducible over K for all $n \geq 1$. Suppose $\mathfrak{p} \subset \mathcal{O}_K$ does not ramify in $K_{n,t} \simeq K[x]/(\Phi_n(x,t))$ for any n. Then, for $k \geq 1$, the number of degree k primes of $K_{n,t}$ lying over $\mathfrak{p}$ is N/k, where N is the number of paths of length n in the graph of ϕ acting on $\overline{\mathbb{F}}_p$ which start with a vertex of weight k and end at $\overline{t}$, the weight 1 vertex corresponding to the image of t in $\mathbb{F}_p$. [The weight of a vertex is the exact degree of the vertex over $\mathbb{F}_p$.]

Date: September 19, 2012.

Key words and phrases. Chebyshev polynomial, iterated polynomial, post-critically finite map, finite field.

The Chebyshev polynomials form a very interesting family of postcritically finite functions. In section 3, we combine our results with those of [1] to give a rule describing how certain primes split in the tower of number fields generated by adjoining the roots of $\Phi_{n,\ell}(x,t) := T_\ell^{\circ n}(x) - t$.

Remark 1. The Chebyshev polynomials that we refer to in this paper are a normalized version of the Chebyshev polynomials of the first kind. These are monic polynomials in $\mathbb{Z}[x]$ defined by the relation

$$T_n(2\cos\theta) = 2\cos(n\theta),$$

and also satisfy the recurrence

$$T_0(x) = 2; \quad T_1(x) = x; \quad T_{n+1}(x) = x\,T_n(x) - T_{n-1}(x).$$

Other properties of these polynomials are presented in the appendix. Also see [2] for a deeper description of these polynomials.

Notation 2. Throughout this pager, we use $\nu_\ell\colon \mathbb{Z} \to \mathbb{Z}$ to denote the ℓ-adic valuation

2. Description of Graphs

For the rest of this paper we assume that ℓ is a prime number, p is an odd prime different from ℓ, and n is a positive integer. Let $G(\ell, p, n)$ denote the graph of T_ℓ acting on $\mathbb{F}_{p^n}$. The analysis we provide in this paper can also be applied to the cases where $p = 2$ and $p = \ell$. In most instances, only minor adjustments in our arguments are needed to address these special cases. Furthermore, although we only consider the Chebyshev polynomials of prime degree, the composition property $T_{mn}(x) = T_m(T_n(x))$ would allow one to extend the results from the prime degree case to Chebyshev polynomials of arbitrary degree.

Given $a \in \mathbb{F}_{p^n}$, define the polynomial $u(x) = x^2 - ax + 1$. Let $\alpha \in \mathbb{F}_{p^{2n}}^\times$ be a root of this polynomial.

Lemma 2.1. For any $k \geq 0$, $T_k(a) = \alpha^k + \alpha^{-k}$.

Proof. We proceed by induction on k. For the base cases, we have

$$T_0(a) = 2 = \alpha^0 + \alpha^0 \quad \text{and} \quad T_1(a) = a = \alpha + \alpha^{-1}.$$

Now assume that $T_k(a) = \alpha^k + \alpha^{-k}$ for all positive integers up to k. The recursion relation of the Chebyshev polynomials gives

$$T_{k+1}(a) = a\,T_k(a) - T_{k-1}(a) = (\alpha + \alpha^{-1})(\alpha^k + \alpha^{-k}) - (\alpha^{k-1} + \alpha^{-(k-1)}) = \alpha^{k+1} + \alpha^{-(k+1)}.$$

□

Since we are working over finite fields, every element $a \in \mathbb{F}_{p^n}$ is necessarily preperiodic under the action of T_ℓ. The graph of the forward orbit of a is a string of points of length t leading into a cycle of length c where t is the preperiod of a and c is the period of $T_\ell^{\circ t}(a)$. (See figure 1.)

The following theorem provides a method for calculating the preperiod of a as well as the length of the cycle to which a is attached.

Theorem 2.2. Suppose $a \in \mathbb{F}_{p^n}$ has preperiod t, and $T_\ell^{\circ t}(a)$ has period c. Let $\alpha \in \mathbb{F}_{p^{2n}}^\times$ be a root of $x^2 - ax + 1$, and write $\operatorname{ord}_{\mathbb{F}_{p^{2n}}^\times} \alpha = \ell^\lambda \omega$, where $\gcd(\omega, \ell) = 1$. Then

$$t = \lambda, \quad \text{and} \quad c = \operatorname{ord}_{(\mathbb{Z}/\omega\mathbb{Z})^\times/(\pm 1)} \ell,$$

i.e. λ is the ℓ-valuation of the order of α in $\mathbb{F}_{p^{2n}}^\times$, and c is the least positive integer such that $\ell^{2c} \equiv 1 \pmod{\omega}$.



FIGURE 1. The graph of the forward orbit of a.

Proof. The element a satisfies the relation $T_\ell^{\circ\, t+c}(a) = T_\ell^{\circ\, t}(a)$, which leads to the following equivalences.

$$\begin{aligned} T_\ell^{\circ\, t+c}(a) = T_\ell^{\circ\, t}(a) \quad &\Leftrightarrow \quad \alpha^{\ell^{t+c}} + \alpha^{-\ell^{t+c}} = \alpha^{\ell^t} + \alpha^{-\ell^t} \\ &\Leftrightarrow \quad \alpha^{\ell^{t+c}}\alpha^{\ell^{t+c}} - \alpha^{\ell^t}\alpha^{\ell^{t+c}} - \alpha^{-\ell^t}\alpha^{\ell^{t+c}} + 1 = 0 \\ &\Leftrightarrow \quad \left(\alpha^{\ell^{t+c}} - \alpha^{\ell^t}\right)\left(\alpha^{\ell^{t+c}} - \alpha^{-\ell^t}\right) = 0 \\ &\Leftrightarrow \quad \alpha^{\ell^{t+c}} = \alpha^{\ell^t} \quad \text{or} \quad \alpha^{\ell^{t+c}} = \alpha^{-\ell^t} \\ &\Leftrightarrow \quad \alpha^{\ell^t(\ell^c-1)} = 1 \quad \text{or} \quad \alpha^{\ell^t(\ell^c+1)} = 1 \\ &\Leftrightarrow \quad \ell^\lambda\omega \mid \ell^t(\ell^c-1) \quad \text{or} \quad \ell^\lambda\omega \mid \ell^t(\ell^c+1) \end{aligned}$$

which implies that $\lambda \mid t$ and $\ell^c \equiv \pm 1 \pmod{\omega}$.

Note that λ cannot be strictly less than t, otherwise the equivalences above imply that $T_\ell^{\circ\,\lambda+c}(a) = T_\ell^{\circ\,\lambda}(a)$, contradicting the minimality of t.

Similarly, if there existed c' strictly less than c such that $\ell^{c'} \equiv \pm 1 \pmod{\omega}$, then $\alpha^{\ell^t(\ell^{c'}\pm 1)} = 1$ up to a choice of sign. This would imply that a satisfies the relation $T_\ell^{\circ\, t+c'}(a) = T_\ell^{\circ\, t}(a)$, contradicting the minimality of c. ☐

The possible orders for α can be described completely.

Theorem 2.3. (1) For $a = 2$, we have $\alpha = 1$ and $\operatorname{ord}_{\mathbb{F}_{p^{2n}}^\times} \alpha = 1$.

(2) For $a = -2$, we have $\alpha = -1$ and $\operatorname{ord}_{\mathbb{F}_{p^{2n}}^\times} \alpha = 2$.

(3) For each divisor d of $p^n - 1$, $d \neq 1, 2$, there are $\varphi(d)/2$ elements $a \in \mathbb{F}_{p^n}$ for which the corresponding α has $\operatorname{ord}_{\mathbb{F}_{p^{2n}}^\times} \alpha = d$.

(4) For each divisor d of $p^n + 1$, $d \neq 1, 2$, there are $\varphi(d)/2$ elements $a \in \mathbb{F}_{p^n}$ for which the corresponding α has $\operatorname{ord}_{\mathbb{F}_{p^{2n}}^\times} \alpha = d$

Proof. Let $a \in \mathbb{F}_{p^n}$ and define $u(x) = x^2 - ax + 1$. If u is reducible over $\mathbb{F}_{p^n}$, then u has roots α and α^{-1} in $\mathbb{F}_{p^n}^\times$. This unit group is a cyclic group of order $p^n - 1$ and therefore contains one element of order 1, one element of order 2, and $\varphi(d)$ elements of order d for every divisor $d > 2$ of $p^n - 1$. The element of order one, $\alpha = 1$, corresponds to $a = 2$, and the element of order two, $\alpha = -1$, corresponds to $a = -2$. For all other divisors $d > 2$, α and α^{-1} are distinct, and therefore each pair $\{\alpha, \alpha^{-1}\}$ corresponds to a single element a. This completes cases (1), (2), and (3).

If u is irreducible over $\mathbb{F}_{p^n}$, then u has roots α and α^{-1} in $\mathbb{F}_{p^{2n}}^\times$. In particular, $\alpha \notin \mathbb{F}_{p^n}^\times$, so the Frobenius automorphism gives $\alpha^{p^n} = \alpha^{-1}$. Therefore α is a $p^n + 1$ root of unity, as it satisfies $\alpha^{p^n+1} = \alpha^{-1}\alpha = 1$. The argument used above now applies, and we conclude (4). ☐

We now describe a method for identifying all the points with a given preperiod.

Theorem 2.4. Let ν_ℓ denote the ℓ-adic valuation, let δ be a generator for $\mathbb{F}_{p^{2n}}^\times$, and define

$$\theta = \delta^{p^n-1} \quad \text{and} \quad \gamma = \delta^{p^n+1}.$$

The periodic points in $G(\ell, p, n)$ are

$$\{\theta^i + \theta^{-i} \colon 1 \le i \le (p^n-1)/2 \text{ and } \nu_\ell(i) \ge \nu_\ell(p^n+1)\}$$
$$\cup \{\gamma^j + \gamma^{-j} \colon 0 \le j \le (p^n-1)/2 \text{ and } \nu_\ell(j) \ge \nu_\ell(p^n-1)\},$$

and the points with preperiod $t \ge 1$ are

$$\{\theta^i + \theta^{-i} \colon 1 \le i \le (p^n-1)/2 \text{ and } \nu_\ell(i) = \nu_\ell(p^n+1) - t\}$$
$$\cup \{\gamma^j + \gamma^{-j} \colon 0 \le j \le (p^n-1)/2 \text{ and } \nu_\ell(j) = \nu_\ell(p^n-1) - t\}.$$

Proof. For $a \in \mathbb{F}_{p^n}$, let $u(x) = x^2 - ax + 1$ with roots α and $\alpha^{-1} \in \mathbb{F}_{p^{2n}}^\times$. We consider the two cases where u is irreducible and u is reducible over $\mathbb{F}_{p^n}$.

Suppose that u is irreducible over $\mathbb{F}_{p^n}$. Then $\alpha = \theta^i$ for some i where $1 \le i \le (p^n-1)/2$. Write $p^n + 1 = \ell^\lambda \omega$ where $\gcd(\omega, \ell) = 1$. By Theorem 2.2, a is periodic if and only if there exists c such that $\alpha^{\ell^c-1} = 1$ or $\alpha^{\ell^c+1} = 1$. That is, a is periodic if and only if there exists c such that $\theta^{i(\ell^c \pm 1)} = 1$, if and only if there exists c such that $p^n + 1 = \ell^\lambda\omega \mid i(\ell^c \pm 1)$, if and only if $\ell^\lambda \mid i$ and there exists c such that $\omega \mid \ell^c \pm 1$. We know such c exist; for instance take $c = \operatorname{ord}_{(\mathbb{Z}/\omega\mathbb{Z})^\times/(\pm 1)} \ell$. Thus a is periodic if and only if $\nu_\ell(i) \ge \nu_\ell(p^n+1)$.

Assuming still that u is irreducible, we take on the case where a has preperiod $t \ge 1$. Theorem 2.2 tells us that a is preperiodic with preperiod $t \ge 1$ if and only if there exists c such that $\theta^{i\ell^t(\ell^c \pm 1)} = 1$ and $\theta^{i\ell^{t-1}(\ell^c \pm 1)} \ne 1$. Following the same argument as above, we see that a is preperiodic with preperiod t if and only if $\ell^\lambda \mid i\ell^t$ and $\ell^\lambda \nmid i\ell^{t-1}$. Thus a has preperiod t if and only if $\nu_\ell(i) = \nu_\ell(p^n+1) - t$.

The case where u is reducible over $\mathbb{F}_{p^n}$ follows a similar argument where instead $\alpha = \gamma^j$ for some j where $0 \le j \le (p^n-1)/2$, and the order of α is divisible by $p^n - 1$. ☐

We can now make an exact count of the number of points in $G(\ell, p, n)$ at every preperiod.

Notation 3. We write $p^n - 1 = \ell^{\lambda_1}\omega_1$ and $p^n + 1 = \ell^{\lambda_2}\omega_2$ where ω_1 and ω_2 are relatively prime to ℓ. We define $m \in \{1, 2\}$ to be the index such that $\lambda_m = \max\{\lambda_1, \lambda_2\}$.

Theorem 2.5. (1) If $\ell = 2$, the number of points with preperiod t in $G(2, p, n)$ is

Preperiod	$t = 0$	$t = 1$	$2 \le t \le \lambda_m$
# of points	$\frac{\omega_1 + \omega_2}{2}$	$\frac{\omega_1 + \omega_2}{2}$	$2^{t-2}\,\omega_m$

(2) If ℓ is odd, the number of points with perperiod t in $G(\ell, p, n)$ is

Preperiod	$t = 0$	$1 \le t \le \lambda_m$
# of points	$\frac{\omega_1 + \omega_2}{2}$	$(\ell^t - \ell^{t-1})\frac{\omega_m}{2}$

Proof. We begin with the case where $\ell = 2$.

Periodic points (t=0): By theorem 2.4, the number of periodic points in $G(2,p,n)$ is

$$\begin{aligned}&\#\{1 \le i \le (p^n-1)/2\colon \nu_2(i) \ge \nu_2(p^n+1)\}\\&\qquad + \#\{0 \le j \le (p^n-1)/2\colon \nu_2(j) \ge \nu_2(p^n-1)\}.\end{aligned}$$

The size of the first set is

$$\begin{aligned}&\#\{1 \le i \le (p^n-1)/2\colon \nu_2(i) \ge \nu_2(p^n+1)\}\\&\quad = \#\{i \in \mathbb{Z}\colon 1 \le 2^{\lambda_2} i \le \frac{p^n-1}{2} = \frac{p^n+1}{2} - 1 = 2^{\lambda_2-1}\omega_2 - 1\}\\&\quad = \#\{i \in \mathbb{Z}\colon 1 \le 2i \le \omega_2 - 1\}\\&\quad = \frac{\omega_2-1}{2}\end{aligned}$$

The size of the second set is

$$\begin{aligned}&\#\{0 \le j \le (p^n-1)/2\colon \nu_2(j) \ge \nu_2(p^n-1)\}\\&\quad = \#\{j \in \mathbb{Z}\colon 0 \le 2^{\lambda_1} j \le (p^n-1)/2 = 2^{\lambda_1-1}\omega_1\}\\&\quad = \#\{j \in \mathbb{Z}\colon 0 \le 2j \le \omega_1\}\\&\quad = \frac{\omega_1+1}{2}.\end{aligned}$$

Thus, the total number of periodic points is

$$\frac{\omega_1+1}{2} + \frac{\omega_2-1}{2} = \frac{\omega_1+\omega_2}{2}.$$

Preperiod t=1: By theorem 2.4, the number of points with perperiod 1 is

$$\begin{aligned}&\#\{1 \le i \le (p^n-1)/2\colon \nu_2(i) = \nu_2(p^n+1) - 1\}\\&\qquad + \#\{0 \le j \le (p^n-1)/2\colon \nu_2(j) = \nu_2(p^n-1) - 1\}.\end{aligned}$$

The size of the first set is

$$\begin{aligned}&\#\{1 \le i \le (p^n-1)/2\colon \nu_2(i) = \nu_2(p^n+1) - 1\}\\&\quad = \#\{i \in \mathbb{Z}\colon 1 \le 2^{\lambda_2-1} i \le \frac{p^n+1}{2} - 1 = 2^{\lambda_2-1}\omega_2 - 1, \gcd(i,2) = 1\}\\&\quad = \#\{i \in \mathbb{Z}\colon 1 \le i \le \omega_2 - 1, \gcd(i,2) = 1\}\\&\quad = \frac{\omega_2-1}{2}.\end{aligned}$$

The size of the second set is

$$\begin{aligned}&\#\{0 \le j \le (p^n-1)/2\colon \nu_2(j) = \nu_2(p^n-1) - 1\}\\&\quad = \#\{j \in \mathbb{Z}\colon 0 \le 2^{\lambda_1-1} j \le (p^n-1)/2 = 2^{\lambda_1-1}\omega_1, \gcd(j,2) = 1\}\\&\quad = \#\{j \in \mathbb{Z}\colon 0 \le j \le \omega_1, \gcd(j,2) = 1\}\\&\quad = \frac{\omega_1+1}{2}.\end{aligned}$$

Thus the number of points with preperiod 1 is

$$\frac{\omega_1+1}{2} + \frac{\omega_2-1}{2} = \frac{\omega_1+\omega_2}{2}.$$

Preperiod $2 \leq t \leq \lambda_m$*:* By theorem 2.4, the number of points with perperiod $t \geq 2$ is

$$\begin{aligned}&\#\{1 \leq i \leq (p^n-1)/2\colon \nu_2(i) = \lambda_2 - t\}\\ &\qquad + \#\{0 \leq j \leq (p^n-1)/2\colon \nu_2(j) = \lambda_1 - t\}.\end{aligned}$$

One of these sets is necessarily empty since $p^n - 1$ and $p^n + 1$ are consecutive even integers, and therefore $\min\{\lambda_1, \lambda_2\} = 1$. The size of the nonempty set is equal to

$$\begin{aligned}&\#\{i \in \mathbb{Z}\colon 0 \leq 2^{\lambda_m - t} i \leq 2^{\lambda_m - 1}\omega_m, \gcd(i,2) = 1\}\\ &\quad = \#\{i \in \mathbb{Z}\colon 0 \leq i \leq 2^{t-1}\omega_m, \gcd(i,2) = 1\}\\ &\quad = 2^{t-2}\omega_m.\end{aligned}$$

The argument for $\ell > 2$ is essentially the same but with fewer cases to consider since ℓ divides at most one of $p^n + 1$ and $p^n - 1$. If either of these values is divisible by ℓ, then λ_m is positive and $G(\ell, p, n)$ contains non-periodic points. Otherwise $\lambda_1 = \lambda_2 = 0$ and all points are strictly periodic. □

The connected components of these graphs have two distinguishing features: 1) a unique cycle comprised of the periodic points, and 2) a set of paths leading into the cycle containing the non-periodic points. Since all points are preperiodic, every connected component necessarily contains at least one periodic point, and therefore contains a cycle. The components may or may not contain non-periodic points, but when they do, the non-periodic points have a very specific structure. To describe the arrangement of the non-periodic points, we introduce some notation from graph theory.

We say that a connected graph G is a tree if it contains no cycles. A tree is called rooted if one vertex is specified as a root. Designating a root imparts an orientation on the graph whereby all edges are directed towards the root. In a rooted tree, the parent u of a vertex v is the unique vertex connected to v on the path connecting v to the root, and v is said to be a child of u. An n-ary tree is a rooted tree in which each parent has at most n children, and we say that a tree is complete n-ary if each parent has exactly n children. The height of a vertex v is the length of the path connecting v to the root, and the height of the tree is the maximal height taken over all vertices, if it exists.

The following theorem describes the structure of all connected components of the graph by determining the lengths of cycles, the number of cycles of each length, and the arrangement of non-periodic points attached to each cycle.

Theorem 2.6. For each divisor $d > 2$ of ω_1, let $c = \operatorname{ord}_{(\mathbb{Z}/d\mathbb{Z})^\times/(\pm 1)} \ell$. Then $G(\ell, p, n)$ contains $\varphi(d)/(2c)$ cycles of length c. Attached to each point in the cycles are $\ell - 1$ complete ℓ-ary rooted trees of height $\lambda_1 - 1$, provided $\lambda_1 \geq 1$.

Similarly, for each divisor $d > 2$ of ω_2, let $c = \operatorname{ord}_{(\mathbb{Z}/d\mathbb{Z})^\times/(\pm 1)} \ell$. Then $G(\ell, p, n)$ contains $\varphi(d)/(2c)$ cycles of length c. Attached to each point in the cycles are $\ell - 1$ complete ℓ-ary rooted trees of height $\lambda_2 - 1$, provided $\lambda_2 \geq 1$.

If ℓ is odd, then the points 2 and -2 are fixed, and from each of these points hang $(\ell - 1)/2$ complete ℓ-ary rooted trees of height $\lambda_m - 1$, provided $\lambda_m \geq 1$.

If $\ell = 2$, then the graph contains the edges $(0 \to -2), (-2 \to 2), (2 \to 2)$, and 0 is the root of a complete binary tree of height $\lambda_m - 2$.

Proof. Let $d > 2$ be a divisor of ω_1. By theorems 2.3 and 2.2, there are $\varphi(d)/2$ periodic elements of $\mathbb{F}_{p^n}$ of period c. Furthermore, this set is fixed by T_ℓ, for if α is an element of order d in $\mathbb{F}_{p^{2n}}^\times$, then α^ℓ is also an element of order d since ℓ and d are relatively prime. Thus the $\varphi(d)/2$ elements are arranged in $\varphi(d)/(2c)$ cycles of length c.

The points with perperiod t that lead into these cycles are exactly the points whose corresponding α have order $d\ell^t$. Thus there are $\varphi(d\ell^t)/2 = \ell^{t-1}(\ell-1)\varphi(d)/2$ elements with perperiod t leading into these cycles, provided that $d\ell^t$ divides $p^n - 1$, i.e. $\lambda_1 \geq 1$. Completeness follows from the fact that the number of preimages of any point is bounded by the degree of T_ℓ. The same argument applies to the other three cases. ☐

We now provide a couple of examples (Figures 2 and 3) so that the reader can get a feel for these graphs. To summarize the results of the previous theorems, we recall that each divisor $d > 2$ of $p^n - 1$ and $p^n + 1$ corresponds to $\varphi(d)/2$ distinct points in $\mathbb{F}_{p^n}$. The divisors $d = 1$ and $d = 2$ correspond to the points 2, respectively, -2 in $\mathbb{F}_{p^n}$. If d is relatively prime to ℓ, then the points corresponding to d are periodic. Otherwise, the points corresponding to d are preperiodic with preperiod $\nu_\ell(d)$. The weight of a point is the exact degree of the point over $\mathbb{F}_p$. The double arrows in the graphs denote double roots, and to avoid clutter, only selected vertices have been labeled.

The graph $T_2(x) = x^2 - 2$ acting on $\mathbb{F}_{3^4}$.

$\ell = 2, p = 3, n = 4$					
Divisors of 80	Number of points in $\mathbb{F}_{3^4}$	Period	Preperiod	Weight	Cycles of this type
1	1	1	0	1	1
2	1	-	1	1	
4	1	-	2	1	
8	2	-	3	2	
16	4	-	4	4	
5	2	2	0	2	1
10	2	-	1	2	
20	4	-	2	4	
40	8	-	3	4	
80	16	-	4	4	
Divisors of 82					
41	20	20	0	4	1
82	20	-	1	4	



FIGURE 2. $G(2, 3, 4)$.

Graph of $T_3(x) = x^3 - 3x$ acting on $\mathbb{F}_{53}$.

$\ell = 3, p = 53, n = 1$				
Divisors of 52	Number of points in $\mathbb{F}_{53}$	Period	Preperiod	Cycles of this type
4	1	1	0	1
13	6	3	0	2
26	6	3	0	2
52	12	6	0	2
Divisors 54				
1	1	1	0	1
3	1	-	1	
9	3	-	2	
27	9	-	3	
2	1	1	0	1
6	1	-	1	
18	3	-	2	
54	9	-	3	



FIGURE 3. $G(3, 53, 1)$.

3. DECOMPOSITION OF PRIMES.

As mentioned at the beginning of this paper, one application of the study of these graphs is to understand how primes decompose in certain extensions. Before taking advantage of Proposition 1.1, we complete some missing details.

Proposition 3.1. For $n \geq 1$, the Chebyshev polynomial T_n is postcritically finite.

Proof. The critical points of T_n are the points where the derivative vanishes. The derivative of T_n satisfies the relation

$$\frac{d}{dx}T_n = nU_{n-1}$$

where U_{n-1} is the normalized, degree $n-1$ Chebyshev polynomial of the second kind. [See appendix for proof.] The Chebyshev polynomials of the second kind are defined by the relation

$$U_{n-1}(2\cos\theta) = \frac{\sin(n\theta)}{\sin\theta}.$$

From here it is clear that the roots of U_{n-1} are

$$x_k = 2\cos\left(\frac{k\pi}{n}\right), \quad k = 1, 2, \ldots, n-1.$$

It is left to show that these critical values are preperiodic for T_n. Evaluating T_n at these values, we see that their forward orbits are finite.

$$\begin{aligned}
&T_n\left(2\cos\left(\frac{k\pi}{n}\right)\right) = 2\cos(k\pi) = (-1)^k\, 2, \quad k = 1, 2, \dots n-1;\\
&T_n(-2) = T_n(2\cos(\pi)) = 2\cos(n\pi) = (-1)^n\, 2\\
&T_n(2) = T_n(2\cos(0)) = 2\cos(0) = 2.
\end{aligned}$$

□

Notation 4. For a prime p, we denote by $\bar{\ }$ reduction modulo p.

Recall that for a fixed prime ℓ and an integer t, we define a family of polynomials $\Phi_{n,\ell}(x,t) = T_\ell^{\circ n}(x) - t$. We are specifically interested in the case where $\Phi_{n,\ell}(x,t)$ is irreducible for every n. Locally, we can address this question of irreducibility modulo a prime p by considering the position of $\bar{t}$ in the graph $G(\ell, p, 1)$.

The following lemma describes how the graph $G(\ell, p, k)$ evolves as k varies. Specifically, it gives a formula for the maximum height, λ, of the trees described in Theorem 2.6.

Lemma 3.2. Let $c = \operatorname{ord}_{(\mathbb{Z}/\ell\mathbb{Z})^\times/(\pm 1)} p$ and $\lambda = \max\{\nu_\ell(p^k-1), \nu_\ell(p^k+1)\}$.

(1) If ℓ is odd, then

$$\lambda = \nu_\ell(p^{2k}-1) = \begin{cases} \nu_\ell(p^{2c}-1) + \nu_\ell(k) \geq 1 & \text{if } c \mid k \\ 0 & \text{otherwise} \end{cases}.$$

(2) If $\ell = 2$, then $c = 1$ and

$$\lambda = \nu_2(p^{2k}-1) - 1 = \nu_2(p^2-1) + \nu_2(k) - 1 \geq 2.$$

Proof. Beginning with ℓ odd, we see that $p^{2k} - 1 = (p^k+1)(p^k-1)$ is divisible by ℓ if and only if $p^k \equiv \pm 1 \pmod{\ell}$, and therefore c divides k. Furthermore, if ℓ divides $p^{2k}-1$, then ℓ divides exactly one of p^k+1 and p^k-1, and we have

$$\nu_\ell(p^{2k}-1) = \nu_\ell(p^k+1) + \nu_\ell(p^k-1) = \lambda \geq 1.$$

Otherwise, if $\ell = 2$, both p^k-1 and p^k+1 are even with $\min\{\nu_2(p^k-1), \nu_2(p^k+1)\} = 1$ and $\lambda \geq 2$, so

$$\nu_\ell(p^{2k}-1) = \nu_\ell(p^k+1) + \nu_\ell(p^k-1) = \lambda + 1 \geq 3.$$

We are now left to show that $\nu_\ell(p^{2k}-1) = \nu_\ell(p^{2c}-1) + \nu_\ell(k)$ provided that c divides k. We begin by writing $k = cm$ for some integer m, and we note that $p^{2c} \equiv 1 \pmod{\ell}$. Then

$$p^{2k} - 1 = p^{2cm} - 1 = (p^{2c})^m - 1 = (p^{2c}-1)\big((p^{2c})^{m-1} + (p^{2c})^{m-2} + (p^{2c})^{m-3} + \cdots + 1\big).$$

Since all powers of p^{2c} are congruent to 1 mod ℓ, we have

$$\nu_\ell(p^{2k}-1) = \nu_\ell(p^{2c}-1) + \nu_\ell(m).$$

Finally, c is a divisor of $\ell - 1$ since it is the order of p in $(\mathbb{Z}/\ell\mathbb{Z})^\times/(\pm 1)$, and in particular, $\nu_\ell(c) = 0$. Thus

$$\nu_\ell(m) = \nu_\ell(c) + \nu_\ell(m) = \nu_\ell(cm) = \nu_\ell(k),$$

completing the proof. □

Proposition 3.3. Suppose $p \equiv \pm 1 \pmod{\ell}$. If $\bar{t}$ is of maximal height in $G(\ell, p, 1)$, then $\Phi_{n,\ell}(x,t)$ is irreducible modulo p.

Proof. Let p be a prime congruent to ± 1 modulo ℓ. Then $c = \mathrm{ord}_{(\mathbb{Z}/\ell\mathbb{Z})^\times/(\pm 1)} p = 1$, so by Lemma 3.2 maximal height attained by the trees in $G(\ell, p, 1)$ is $\nu_\ell(p^2 - 1)$. Similarly, the maximal height attained by the trees in $G(\ell, p, \ell^n)$ (respectively, $G(\ell, p, \ell^{n-1})$) is $\nu_\ell(p^2-1)+n$ (respectively, $\nu_\ell(p^2-1)+n-1$).

Let t be an integer such that $\overline{t}$ is of maximal height in $G(\ell, p, 1)$. That is, $\overline{t}$ is preperiodic with preperiod $\nu_\ell(p^2-1)$ with regard to the action of T_ℓ on $\mathbb{F}_p$. By Theorem 2.6, the tree in $G(\ell, n, \ell^n)$ (respectively, $G(\ell, n, \ell^{n-1})$) rooted at $\overline{t}$ is a complete ℓ-ary tree of height n (respectively, height $n-1$). Note that $\theta \in \mathbb{F}_{p^k}$ is a root of $\overline{\Phi}_{n,\ell}(x,t)$ if any only if there exists a path of length n from θ to $\overline{t}$ in $G(\ell, p, k)$. Since there are no paths of length n terminating at $\overline{t}$ in $G(\ell, p, \ell^{n-1})$ and ℓ^n paths of length n terminating at $\overline{t}$ in $G(\ell, p, \ell^n)$, we see that $\mathbb{F}_{p^{\ell^n}}$ is the smallest field to contain a root of $\overline{\Phi}_{n,\ell}(x,t)$ (in fact, it contains all the roots). In particular, if θ is a root of $\overline{\Phi}_{n,\ell}(x,t)$, then $\mathbb{F}_p(\theta) \simeq \mathbb{F}_{p^{\ell^n}}$ is the degree ℓ^n extension of $\mathbb{F}_p$, and therefore $\overline{\Phi}_{n,\ell}(x,t)$ is the minimal polynomial for θ. ☐

Corollary 3.4. Suppose that ℓ is an odd prime. Then $\Phi_{n,\ell}(x,t)$ is reducible modulo p if and only if $\Phi_{1,\ell}(x,t)$ has a root modulo p.

Proof. Clearly, if $\Phi_{1,\ell}(x,t)$ has a root modulo p then $\Phi_{n,\ell}(x,t)$ is reducible. For if $(x-a)$ is a factor of $T_\ell(x)-t$ for some $a \in \mathbb{F}_p$, then $(T_\ell^{\circ(n-1)} - a)$ is a factor of $\Phi_{n,\ell}(x,t) = T_\ell^{\circ n}(x) - t = T_\ell(T_\ell^{\circ(n-1)}(x)) - t$.

On the other hand, Proposition 3.3 implies that if $\Phi_{n,\ell}(x,t)$ is reducible modulo p, then $\overline{t}$ does not achieve maximal height in $G(\ell, p, 1)$, and thus $\Phi_{1,\ell}(x,t)$ has a root modulo p. ☐

We can now establish that for a fixed ℓ, there are infinitely many choices for t such that $\Phi_{n,\ell}(x,t)$ is a family of irreducible polynomials. Our condition for irreducibility relies on the existence of a prime p such that $p \equiv \pm 1 \pmod{\ell}$. However, it is known that infinitely many such primes exist thanks to Dirichlet's theorem on arithmetic progressions. For any of these primes p, we construct the graph $G(\ell, p, 1)$ and identify a $\overline{t}$ of maximal preperiod. Then, for any integer t such that $t \equiv \overline{t} \pmod{p}$, $\Phi_{n,\ell}(x,t)$ is irreducible for all n.

We now proceed to show that only finitely many primes ramify in the tower of fields

$$\mathbb{Q} = K_{0,t} \subset K_{1,t} \subset K_{2,t} \subset \cdots, \quad K_{n,t} \simeq K[x]/(\Phi_{n,\ell}(x,t))$$

by showing that the primes dividing the discriminants of the polynomials $\Phi_{n,\ell}(x,t)$ is a finite set. Since the number field discriminant divides the polynomial discriminant, only finitely many primes divide the discriminants of the fields $K_{n,t}$, and therefore only finitely many primes ramify in the tower of fields. The following proposition gives a formula for the discriminant for an iterated polynomial, and from this we derive a formula for the discriminant of $\Phi_{n,\ell}(x,t)$.

Notation 5. Let K be a number field and $\phi = a_d x^d + \cdots + a_0 \in K[x]$. We identify the multiplicities of the critical points of ϕ by

$$\phi'(x) = d a_d \prod_{r \in \mathcal{R}_\phi} (x - r)^{m_r(\phi)},$$

where $\mathcal{R}_\phi$ is the set of critical points of ϕ. For each critical value $\beta \in \mathcal{B}_\phi$, define

$$\mathcal{M}_\beta(\phi) = \sum_{r \in \mathcal{R}_\phi,\, \phi(r) = \beta} m_r(\phi).$$

Proposition 3.5. [Aitken, Hajir, Maire] For $\phi = a_d x^d + \cdots + a_0 \in K[x]$ and $n \geq 1$

$$\mathrm{disc}_x(\Phi_n) = (-1)^{(d^n-1)(d^n-2)/2} d^{nd^n} a_d^{(d^n-1)^2/(d-1)} \prod_{\beta \in \mathcal{B}_{\phi^{\circ n}}} (t - \beta)^{\mathcal{M}_\beta(\phi^{\circ n})}$$

where $\Phi_n(x,t) = \phi^{\circ n}(x) - t$.

Corollary 3.6. Let ℓ be a prime and define $\Phi_{n,\ell}(x,t) = T_\ell^{\circ n}(x) - t$ as before.

(1) If ℓ is odd, then
$$\operatorname{disc}_x(\Phi_{n,\ell}(x,t)) = (-1)^{(\ell^n-1)/2}\ell^{n\ell^n}(t^2-4)^{(\ell^n-1)/2},$$
and the discriminants satisfy the recursion
$$\operatorname{disc}_x(\Phi_{n+1,\ell}(x,t)) = (-1)^{(\ell-1)/2}\ell^{\ell^{n+1}}\big(\operatorname{disc}_x(\Phi_{n,\ell}(x,t))\big)^\ell(t^2-4)^{(\ell-1)/2}.$$

(2) If $\ell = 2$, then
$$\operatorname{disc}_x(\Phi_{n,2}(x,t)) = (-1)^{(2^n-1)(2^{n-1}-1)}2^{n2^n}(t+2)^{2^{n-1}}(t-2)^{2^{n-1}-1},$$
and the discriminants satisfy the recursion
$$\operatorname{disc}_x(\Phi_{n+1,2}(x,t)) = 4^{2^n}\big(\operatorname{disc}_x(\Phi_{n,2}(x,t))\big)^2(2-t).$$

Proof. We begin with the case where ℓ is odd. In Proposition 3.1, we identified the $\ell^n - 1$ distinct critical points of $T_\ell^{\circ n}(x)$, and identified 2 and -2 as the two critical values. Each critical value is the image of half of the critical points, so $\mathcal{M}_2(T_\ell^{\circ n}) = \mathcal{M}_{-2}(T_\ell^{\circ n}) = (\ell^n-1)/2$ and we can apply Proposition 3.5.

As for the recursion, we use the fact that
$$\frac{\ell^{n+1}-1}{2} = \frac{\ell^{n+1}-\ell+\ell-1}{2} = \frac{\ell(\ell^n-1)}{2} + \frac{\ell-1}{2}$$
to conclude that
$$\begin{aligned}\operatorname{disc}_x(\Phi_{n+1,\ell}(x,t)) &= (-1)^{(\ell^{n+1}-1)/2}\ell^{(n+1)\ell^{n+1}}(t^2-4)^{(\ell^{n+1}-1)/2}\\ &= (-1)^{(\ell-1)/2}\big((-1)^{(\ell^n-1)/2}\big)^\ell\big(\ell^{n\ell^n}\big)^\ell\ell^{\ell^{n+1}}\big((t^2-4)^{(\ell^n-1)/2}\big)^\ell(t^2-4)^{(\ell-1)/2}\\ &= (-1)^{(\ell-1)/2}\big(\operatorname{disc}_x(\Phi_{n,\ell}(x,t))\big)^\ell(t^2-4)^{(\ell-1)/2}.\end{aligned}$$

In the case $\ell = 2$, $T_2^{\circ n}$ has $2^n - 1$ critical points, and it is a quick check to verify that $\mathcal{M}_2(T_2^{\circ n}) = 2^{n-1} - 1$ and $\mathcal{M}_{-2}(T_2^{\circ n}) = 2^{n-1}$, and the discriminant formula follows immediately. For the recursion, we note that $(2^n-1)(2^{n-1}-1)$ is odd for $n \geq 2$. Thus
$$\begin{aligned}\operatorname{disc}_x(\Phi_{n+1,2}(x,t)) &= -2^{(n+1)2^{n+1}}(t+2)^{2^n}(t-2)^{2^n-1}\\ &= -4^{2^n}\big(2^{n2^n}\big)^2\big((t+2)^{2^{n-1}}\big)^2\big((t-2)^{2^{n-1}-1}\big)^2(t-2)\\ &= 4^{2^n}\big(\operatorname{disc}_x(\Phi_{n,2}(x,t))\big)^2(2-t).\end{aligned}$$
□

Notation 6. Throughout the rest of this section, we will use the notation $G(\ell,p,\ell^\infty)$ to denote the union of the nested graphs
$$G(\ell,p,\ell^0) \subset G(\ell,p,\ell^1) \subset G(\ell,p,\ell^2) \subset \cdots \subset G(\ell,p,\ell^\infty).$$

Example 3.7. Consider the family of polynomials $\Phi_{n,\ell}(x,t) = T_\ell^{\circ n}(x) - t$ and the fields they generate $K_{n,t} \simeq \mathbb{Q}[x]/(\Phi_{n,\ell}(x,t))$ when $\ell = 2$ and $t = 0$. The tower $\mathbb{Q} \subset K_{1,0} \subset K_{2,0} \subset \cdots$ is the cyclotomic $\mathbb{Z}_2$-extension of $\mathbb{Q}$. The splitting of primes is completely described by the cyclotomic reciprocity law. (See [4], Theorem 2.13.) Specifically, $K_{n,0} \subset \mathbb{Q}(\zeta_{2^{n+2}})$ is the field fixed by complex conjugation, so cyclotomic reciprocity implies that an odd prime p will split into $2^n/f$ primes, where f is the order of p in $(\mathbb{Z}/(2^{n+2}\mathbb{Z}))^\times/(\pm 1)$, and in particular, p splits completely in $K_{n,0}$ if



FIGURE 4. $G(2,3,1)$

and only if $p \equiv \pm 1 \pmod{2^{n+2}}$. We recover this law using Proposition 1.1 in concert with knowledge of the graph $G(2, p, 2^\infty)$.

In $\mathbb{F}_3$, $\bar{t} = 0$ is the point of maximal preperiod under iteration by T_2. This is evidenced in the graph $G(2, 3, 1)$ (see Figure 4), where we can see that 0 is the point achieving maximal height. Thus by Proposition 3.3, $\Phi_{n,2}(x, 0)$ is irreducible for every $n \geq 1$. Furthermore, by Corollary 3.6, the discriminant of $T_2^{\circ n}(x)$ is a power of 2, and therefore only 2 can ramify in the extensions $K_{n,0}$. Let p be a prime different from 2. By Proposition 1.1, the number of degree k primes lying over p in $K_{n,0}$ is N/k where N is the number of paths of length n in $G(2, p, 2^\infty)$ originating at a point of weight k and ending at 0.

To count these paths, begin by considering the subgraph of $G(2, p, 1)$ containing 0 and its preimages. Theorem 2.6 mandates that 0 is the root of a complete binary tree of maximal height, specifically $\nu_2(p^2 - 1) - 3$. By Lemma 3.2, for any $n \geq 0$, the height of the tree rooted at 0 in $G(2, p, 2^n)$ is

$$\nu_2(p^2 - 1) - 3 + n.$$

By induction on n, we conclude that the points at height $\nu_2(p^2 - 1) - 3 + n$ have weight 2^n. This uniform growth allows us to easily count the paths terminating at 0 in $G(2, p, 2^\infty)$, and we can conclude the following.



FIGURE 5. Subgraphs of $G(2, p, 2^\infty)$ rooted at 0 where $p \equiv \pm 3 \pmod 8$ [left] and $p \equiv \pm 7 \pmod{16}$ [right].

Proposition 3.8. Let p be an odd prime, and define $H = \nu_2(p^2 - 1) - 3$. The number of paths terminating at 0 in $G(2, p, 2^\infty)$ is as follows. For $1 \leq k \leq H$, there are exactly 2^k paths of length k from points of weight one to 0, and for $k > H$ there are exactly 2^k paths of length k from points of weight 2^{k-H} to 0.

Theorem 3.9. Let p be an odd prime, $H = \nu_2(p^2 - 1) - 3$, and let $K_{n,0} = \mathbb{Q}[x]/(T_2^{\circ n}(x))$ as before. Then p splits completely in each extension $K_{0,0}, \ldots, K_{H,0}$ and is completely inert afterwards.

Proof. The result follows immediately from Proposition 3.8 and Proposition 1.1. □

Remark 2. Our result is in complete agreement with the cyclotomic reciprocity law, as we now show. Letting $H = \nu_2(p^2 - 1) - 3$, then $H + 3 = \nu_2(p + 1) + \nu_2(p - 1)$. Therefore $\max\{\nu_2(p + 1), \nu_2(p - 1)\} = H + 2$, and so $p \equiv \pm 1 \pmod{2^{H+2}}$. By cyclotomic reciprocity, p splits completely in $K_{k,0}$ for $k = 0, \ldots, H$.

Example 3.10. Consider the family of polynomials $\Phi_{n,3}(x, 1) = T_3^{\circ n}(x) - 1$. Note that 1 is a point of maximal height in $G(3, 5, 1)$ (see Figure 6), so Proposition 3.3 implies that this is a family of irreducible polynomials. By Corollary 3.6, the only prime divisor of the discriminants of



FIGURE 6. $G(3, 5, 1)$

$\Phi_{n,3}(x,1)$ is 3, so only 3 can ramify in the fields $K_{n,1} = \mathbb{Q}[x]/(\Phi_{n,3}(x,1))$. Let p be a prime different from 3. Note that $T_3(1) \equiv -2 \pmod{p}$ (or 0 in the case $p = 2$), therefore, by Theorem 2.6, 1 is the root of a complete trinary tree of height $\nu_3(p^2-1)-1$. Applying the same analysis as in the previous example, we conclude the following.

Proposition 3.11. Let p be a prime different from 3, and define $H = \nu_3(p^2-1)-1$. The number of paths terminating at 1 in $G(3,p,3^\infty)$ is as follows. For $1 \leq k \leq H$, there are exactly 3^k paths from points of weight one to 1. For $k > H$ there are exactly 3^k paths of length k from points of weight 3^{k-H} to 1.

Theorem 3.12. Let p be a prime different from 3, $H = \nu_3(p^2-1)-1$, and $K_{n,1} = \mathbb{Q}[x]/(T_3^{\circ n}(x)-1)$. Then p splits completely in $\mathbb{Q} = K_{0,1}, \ldots, K_{H,1}$, and is totally inert afterwards.

Proof. The result follows immediately from Proposition 3.11 and Proposition 1.1. □

Remark 3. Here, we have recovered the cyclotomic reciprocity law for the cyclotomic $\mathbb{Z}_3$-extension of $\mathbb{Q}$. In fact, for any odd prime ℓ, we can recover the cyclotomic reciprocity law by considering the polynomials $\Phi_{n,\ell}(x,1)$. We also note that the height H in each of the previous examples is the height of the tree rooted at 0 (respectively, 1) in $G(2,p,1)$ (respectively, $G(3,p,1)$), and is the key value needed to determine the nature of the splitting of the prime. It is, however, not sufficient to only know the value of H. The position of $t \pmod{p}$ also plays a significant role, which we summarize in the following theorem and later illustrate with an example.

Theorem 3.13. Let t be an integer such that $\Phi_{n,\ell}(x,t)$ is a family of irreducible polynomials. Let p be a prime that does not divide the discriminant of $\Phi_{1,\ell}(x,t)$, and denote by $\bar{t}$ the reduction of t modulo p. Let $\alpha \in \mathbb{F}_{p^2}^\times$ be a root of $x^2 - \bar{t}x + 1$ modulo p, and $\lambda = \nu_\ell(\operatorname{ord}_{\mathbb{F}_{p^2}^\times} \alpha)$. Finally, let $H = \nu_\ell(p^2-1) - \lambda$.

(1) If $\lambda = 0$, then there exists at least one prime of degree 1 above p in each field $K_{n,t}$, and exactly one prime of degree 1 above p in $K_{\infty,t} = \varprojlim K_{n,t}$.
(2) If $\lambda \geq 1$ and $\ell \neq 2$, then p splits completely in each field $K_{0,t}, \ldots, K_{H,t}$ and is totally inert afterwards.
(3) If $\lambda \geq 1$, $\ell = 2$, and $\bar{t}$ is contained in a tree of maximal height in $G(\ell,p,1)$, then p splits completely in each field $K_{0,t}, \ldots, K_{H,t}$. If $\bar{t}$ is not contained in a tree of maximal height in $G(\ell,p,1)$, then p is inert in $K_{1,t}$, splits completely in each field $K_{2,t}, \ldots K_{H,t}$, and is completely inert afterwards.

Proof. (1) If $\bar{t}$ is periodic, then there is at least one path of length n from a point of weight one to $\bar{t}$ for every positive integer n. Thus by Proposition 1.1 each field $K_{n,t}$ contains a prime of degree one over p.

(2) If $\lambda \geq 1$ and $\ell \neq 2$, then by Theorem 2.3, $\bar{t}$ is preperiodic with perperiod λ, so the tree in $G(\ell,p,1)$ rooted at $\bar{t}$ has height H. Together, Lemma 3.2 and Theorem 2.6 imply that there are ℓ^n paths of length n in $G(\ell,p,\ell^\infty)$ terminating at $\bar{t}$ all originating at points of weight ℓ^{n-H} (respectively, points of weight 1) if $n > H$ (respectively, if $1 \leq n \leq H$). Applying Proposition 1.1, we achieve our result.

(3) If $\ell = 2$ and $\bar{t}$ is contained in a tree of maximal height in $G(2,p,1)$, then the argument from (2) applies. Otherwise, if $\ell = 2$ and $\bar{t}$ is contained in a tree of minimal height in $G(2,p,1)$, then necessarily $\bar{t}$ is preperiodic with perperiod one. Consider then the tree rooted at $\bar{t}$ in $G(2,p,2^\infty)$. Together, Lemma 3.2 and Theorem 2.6 imply that there are 2^n paths of length n terminating at $\bar{t}$ all originating from points of weight 2^{n-H+1} (respectively, points of weight 2) if $n > H$ (respectively, if $1 \leq n \leq H$). The result follows from Proposition 1.1. □

Example 3.14. Consider the family of polynomials $\Phi_{n,2}(x,105)$ and the tower of fields $K_{n,105} = \mathbb{Q}[x]/(\Phi_{n,2}(x,105))$. These polynomials are irreducible since $105 \equiv 0 \pmod 3$, and 0 is a point of maximal height in $G(2,3,1)$ (see Figure 4). The discriminant formula in Corollary 3.6 shows that only 2, 103, and 107 can ramify in the tower of fields. As before, the splitting of a prime p (different from 2, 103, and 107) in these extensions in completely determined by the tree rooted at $105 \pmod p \in G(2,p,2^\infty)$. Noting that 105 is divisible by 3, 5, and 7, we see that the behavior of 3, 5, and 7 is determined by the tree rooted at 0, and thus the degree at which these primes split is exactly the same as in the cyclotomic $\mathbb{Z}_2$-extension. Namely, 3 and 5 are inert, and 7 splits in $K_{1,2}(x,105)$.

Moving up the list of primes to $p = 11$, we see that $105 \equiv -5 \pmod{11}$, and therefore we must consider the tree rooted at -5 in $G(2,11,2^\infty)$. However, 0 and -5 occupy the same position in the graph $G(2,11,1)$ (see Figure 7), and Lemma 3.2 implies that the tree rooted at -5 is identical to the tree rooted at 0 in $G(2,11,2^n)$ for every n. Thus the splitting behavior of 11 in this extension is also identical to its behavior in the $\mathbb{Z}_2$-extension, that is, it is inert.



FIGURE 7. $G(2,11,1)$

The prime 13 is the smallest prime that displays a different behavior in this tower of fields than in the cyclotomic extension. Since $105 \equiv 1 \pmod{13}$, we focus on the tree rooted at 1 in $G(2,13,2^\infty)$. We can see in $G(2,13,1)$ (see Figure 8) 1 is contained in a tree of maximal height and is the root of a tree of height one. By the previous theorem, 13 splits in $K_{1,105}$, and is otherwise inert.



FIGURE 8. $G(2,13,1)$

4. DENSITY OF PERIODIC POINTS.

Rafe Jones proposed the question: what proportion of points in $G(\ell,p,n)$ are periodic? In particular, does the following limit exist?

$$\lim_{n\to\infty} \frac{\#\{\text{periodic points in } G(\ell,p,n)\}}{p^n}$$

Using our previously established notation, we see that

$$\begin{aligned}
&\lim_{n\to\infty} \frac{\#\{\text{periodic points in } G(\ell,p,n)\}}{p^n} \\
&\quad = \lim_{n\to\infty} \frac{\omega_1+\omega_2}{2p^n} \\
&\quad = \lim_{n\to\infty} \frac{1}{2p^n}\left(\frac{p^n-1}{\ell^{\lambda_1}} + \frac{p^n+1}{\ell^{\lambda_2}}\right) \\
&\quad = \lim_{n\to\infty} \frac{1}{2}\left(\frac{1}{\ell^{\lambda_1}} + \frac{1}{\ell^{\lambda_2}}\right) \\
&\quad = \begin{cases} \lim\limits_{n\to\infty} \dfrac{1}{2\ell^{\lambda_m}} + \dfrac{1}{2} & \text{if } \ell \neq 2 \\ \lim\limits_{n\to\infty} \dfrac{1}{2^{\lambda_m+1}} + \dfrac{1}{4} & \text{if } \ell = 2 \end{cases}.
\end{aligned}$$

Recalling Lemma 3.2, we see that the limits, as written, do not exist since $\lambda_m = 0$ (resp. $\lambda_m = \nu_2(p^2-1)-1$) when ℓ is odd (resp. $\ell = 2$) for infinitely many positive integers n.

This limit, however, can exist if we restrict ourselves to an appropriate tower of fields. For example, consider the sequence

$$\{a_n\}_{n=1}^{\infty} = \{2, \quad 2^2\,3, \quad 2^3\,3^2\,5, \quad 2^4\,3^3\,5^2\,7, \quad \dots \quad , \quad 2^n\,3^{n-1}\,5^{n-2}\,7^{n-3}\cdots p_n, \quad \dots\}$$

and the corresponding tower of fields

$$\mathbb{F}_p \subset \mathbb{F}_{p^{a_1}} \subset \mathbb{F}_{p^{a_2}} \subset \mathbb{F}_{p^{a_3}} \subset \cdots \subset \overline{\mathbb{F}}_p$$

where $\bigcup \mathbb{F}_{p^{a_n}} = \overline{\mathbb{F}}_p$. This sequence of a_n's is constructed so that for any prime ℓ, c divides all but a finitely many a_n, thus $\nu_\ell(a_n)$ (and therefore λ_m) grows monotonically and without bound as n increases. Taking the limit up this tower of fields,

(1) if ℓ is odd,

$$\lim_{n\to\infty} \frac{\#\{\text{periodic points in } G(\ell,p,a_n)\}}{p^{a_n}} = \lim_{n\to\infty} \frac{1}{2\ell^{\nu_\ell(p^{2c}-1)+\nu_\ell(a_n)}} + \frac{1}{2} = \frac{1}{2},$$

(2) if $\ell = 2$,

$$\lim_{n\to\infty} \frac{\#\{\text{periodic points in } G(2,p,a_n)\}}{p^{a_n}} = \lim_{n\to\infty} \frac{1}{2^{\nu_2(p^2-1)+\nu_2(a_n)}} + \frac{1}{4} = \frac{1}{4}.$$

5. Appendix.

In this section we prove some simple relations for the Chebyshev polynomials. We begin by recalling the definition of these polynomials. Again, we remind the reader that the Chebyshev polynomials that we refer to are normalized so as to be monic and integral.

Chebyshev polynomials of the first kind are defined by the relations

$$T_0(x) = 2; \quad T_1(x) = x; \quad T_{n+1}(x) = xT_n(x) - T_{n-1}(x),$$

and also satisfy the relation

$$T_n(2\cos(\theta)) = 2\cos(n\theta).$$

Chebyshev polynomials of the second kind are defined by the relations

$$U_0(x) = 1; \quad U_1(x) = x; \quad U_{n+1}(x) = xU_n(x) - U_{n-1},$$

and also satisfy the relation

$$U_n(2\cos(\theta)) = \frac{\sin((n+1)\theta)}{\sin(\theta)}.$$

Lemma 5.1. The Chebyshev polynomials satisfy the relation

$$T_n(x) = U_n(x) - U_{n-2}(x), \quad n = 2, 3, \dots.$$

Proof. By induction on n. Beginning with the base cases,

$$T_2 = x^2 - 2 = (x^2-1) - 1 = U_2 - U_0;$$
$$T_3 = x^3 - 3x = (x^3 - 2x) - x = U_3 - U_1.$$

Now assume the relation holds for T_n and T_{n-1}. Then

$$\begin{aligned} T_{n+1} = xT_n - T_{n-1} &= x(U_n - U_{n-2}) - (U_{n-1} - U_{n-3}) \\ &= (xU_n - U_{n-1}) - (xU_{n-2} - U_{n-3}) = U_{n+1} - U_{n-1}, \end{aligned}$$

completing the proof. □

Lemma 5.2. The Chebyshev polynomials satisfy the relation

$$\frac{d}{dx}T_n(x) = nU_{n-1}(x), \quad n = 1, 2, \dots.$$

Proof. By induction on n. Beginning with the base cases,

$$\begin{aligned}\frac{d}{dx}T_1 &= \frac{d}{dx}x = 1 = U_1;\\ \frac{d}{dx}T_2 &= \frac{d}{dx}x^2 - 2 = 2x = 2U_2.\end{aligned}$$

Now assume the relation holds for T_n and T_{n-1}. Then

$$\begin{aligned}\frac{d}{dx}T_{n+1} &= \frac{d}{dx}\left(xT_n - T_{n-1}\right)\\ &= xnU_{n-1} + T_n - (n-1)U_{n-2}\\ &= xnU_{n-1} + (U_n - U_{n-2}) - nU_{n-2} + U_{n-2}\\ &= n(xU_{n-1} - U_{n-2}) + U_n = (n+1)U_n,\end{aligned}$$

completing the proof. □

References

[1] W. Aitken, F. Hajir, C. Maire. Finitely Ramified Iterated Extensions. *IMRN*, 14: 855-880, 2005.
[2] J. Silverman, The Arithmetic of Dynamical Systems, Springer, New York, N.Y. 2007.
[3] T. Vasiga, J. Shallit, On the iteration of certain quadratic maps over GF(p), Discrete Mathematics, 277: 219-240, 2004.
[4] L. Washington, Introduction to Cyclotomic Fields, Springer, New York, N.Y. 1997.

E-mail address: `gassert@math.umass.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS, AMHERST, 710 N. PLEASANT STREET, AMHERST, MA, USA 01003



FIGURE 9. A component of $G(3, 67, 9)$.